

**Badminton Return-to-Sport Strategy**  
*Developed by Badminton Canada*

<b>Stage</b>	<b>Aim</b>	<b>Activity</b>	<b>Goal of each step</b>
<b>1</b>	Symptom-limiting activity	Daily activities that do not provoke symptoms	Gradual re-introduction of work/school activities
<b>2</b>	Light aerobic activity	Light jog or stationary cycling at slow to medium pace. No resistance training <i>-Light intensity jogging or stationary cycling for 15-20 minutes at sub-symptom threshold intensity</i>	Increase heart rate
<b>3</b>	Sport-specific exercise	Running <i>- Moderate intensity jogging for 30-60 minutes at sub-symptom threshold intensity</i> <i>- Low to moderate on court activities.</i>	Add movement
<b>4</b>	Non-contact training drills	Harder/longer/more intense on court activity. May start progressive resistance training; <ul style="list-style-type: none"> <li>- <i>Participation in high intensity running and drills</i></li> <li>- <i>Participation in resistance training workouts</i></li> </ul>	Exercise, coordination and increased thinking
<b>5</b>	Full contact practice	Following medical clearance <i>- Participation in full practice without activity restriction</i>	Restore confidence and assess functional skills by coaching staff
<b>6</b>	Return to sport	Normal game play	

Adapted from: McCrory et al. (2017). Consensus statement on concussion in sport – the 5<sup>th</sup> international conference on concussion in sport held in Berlin, October 2016. *British Journal of Sports Medicine*, 51(11), 838-847.